

<u>Tirzepatide 16.6mg/ml Doses</u>	<u>Required units on 1ml syringe</u>
2.5mg	15 units on the syringe
5mg	30 units on the syringe
7.5mg	45 units on the syringe
10mg	60 units on the syringe
12.5mg	75 units on the syringe
15mg	90 units on the syringe
REMEMBER! IF A LOWER DOSE IS WORKING WELL, YOU DO NOT HAVE TO INCREASE YOUR DOSE.	Wells Pharmacy Tirzepatide/Glycine 16.6mg/7.5mg/ml
Note: The FDA and USP require compound pharmacies to recommend multi dose vials only be used for 28 days unless they state the vial can last longer after testing. For the larger vials, you always go with the BUD date listed on the vial. It stays potent much longer than 28 days!	
Notes: This vial has 16.6mg per ml of tirzepatide and 7.5mg per ml of Glycine.	

Dosing rules: Inject subcutaneous once per week in the belly fat region. Start at 2.5mg (15 units) and only increase dose after at least two weeks if the medication is not working and not causing serious side effects. If the medication is curbing appetite and you are losing weight, then stay at your current dose as long as its working! The next doses after 2.5mg (15 units) and in order are 5mg (30 units), 7.5mg (45 units), 10mg (60 units), 12.5mg (75 units) and lastly the max dose of 15mg (90 units). Refer to the dosing schedule to see how many units you need in the syringe to arrive at a specific dose. **AGAIN, ONLY INCREASE DOSE IF THE MEDICATION IS NOT WORKING.**